

Clinical Policy Title:	dalfampridine
Policy Number:	RxA.336
Drug(s) Applied:	Ampyra®
Original Policy Date:	03/06/2020
Last Review Date:	09/14/2020
Line of Business Policy Applies to:	All lines of business

Background

Dalfampridine (Ampyra®) is a potassium channel blocker. Dalfampridine is indicated to improve walking in patients with multiple sclerosis (MS). This was demonstrated by an increase in walking speed.

Dosing Information

Drug Name	Indication	Dosing Regimen	Maximum Dose
Dalfampridine (Ampyra®)	Multiple Sclerosis	10 mg twice daily (approximately 12 hours apart)	20 mg/day

Dosage Forms

- Extended-release tablets: 10 mg

Clinical Policy

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

I. Initial Approval Criteria

A. Multiple Sclerosis (must meet all):

- Diagnosis of MS;
- Prescribed by or in consultation with a neurologist;
- Age 18 years of age or older;
- Member has sustained walking impairment but is able to walk with or without assistance;
- Dose does not exceed 20 mg (2 tablets) per day.

Approval duration

Commercial: 6 months

Medicaid: 6 months

HIM: 6 months

II. Continued Therapy Approval

A. Multiple Sclerosis (must meet all):

- Member is currently receiving dalfampridine that has been authorized by RxAdvance or the member has met initial approval criteria listed in this policy;
- Member is responding positively to therapy;

This clinical policy has been developed to authorize, modify, or determine coverage for individuals with similar conditions. Specific care and treatment may vary depending on individual need and benefits covered by the plan. This policy is not intended to dictate to providers how to practice medicine, nor does it constitute a contract or guarantee regarding payment or results. This document may contain prescription brand name drugs that are trademarks of pharmaceutical manufacturers that are not affiliated with RxAdvance.

3. If request is for a dose increase, new dose does not exceed 20 mg (2 tablets) per day.

Approval duration

Commercial: 12 months

Medicaid: 12 months

HIM: 12 months

III. Appendices

APPENDIX A: Abbreviation/Acronym Key

CrCl: Creatinine clearance

FDA: Food and Drug Administration

MS: Multiple sclerosis

APPENDIX B: Therapeutic Alternatives

None

APPENDIX C: Contraindications/Boxed Warnings

- Contraindication(s):
 - History of seizure; moderate or severe renal impairment (CrCl ≤ 50mL/min); history of hypersensitivity to Ampyra or 4-aminopyridine
- Boxed warning(s):
 - None

Appendix D: General Information

- Use of doses above 10 mg twice daily may increase the risk of seizures.
- Patients with mild renal impairment (CrCl 51-80 mL/min) may exhibit Ampyra levels that approach those attained at higher doses and that have been associated with a higher risk of seizures. Ampyra should be used with caution in this patient population, and CrCl should be estimated or known prior to initiating Ampyra therapy.
- CrCl can be estimated using the Cockcroft-Gault formula: $CrCl = [(140 - age) \times (weight\ in\ kg) \times (0.85\ if\ female)] / (72 \times Cr)$.

References

1. Ampyra Prescribing Information. Ardsley NY: Acorda Therapeutics, Inc; December 2019. Available at <http://www.ampyra.com>. Accessed June 25, 2020.
2. Samkoff LM, Goodman AD. Symptomatic management in multiple sclerosis. *Neurol Clin.* 2011; 29: 449-463.

Review/Revision History	Review/Revised Date	P&T Approval Date
Policy established.	01/2020	03/06/2020
Policy was reviewed: 1) Policy title table was updated 2) Dosing information updated 3) Continued Therapy criteria II.A.1. was rephrased to "Member is currently receiving dalfampridine that has been	07/10/2020	

<p>authorized by RxAdvance or the member has met initial approval criteria listed in this policy”</p> <ul style="list-style-type: none">4) Length of duration for initial therapy and continued therapy approval was updated5) References were updated		
---	--	--