

Clinical Policy Title:	histrelin acetate
Policy Number:	RxA.296
Drug(s) Applied:	Vantas®, Supprelin® LA
Original Policy Date:	02/07/2020
Last Review Date:	06/10/2021
Line of Business Policy Applies to:	All lines of business

Background

Histrelin acetate (Vantas® and Supprelin® LA) is a gonadotropin-releasing hormone (GnRH) agonist.

- Vantas® is indicated for the palliative treatment of advanced prostate cancer.
- Supprelin® LA is indicated for the treatment of children with central precocious puberty (CPP).

Dosing Information

Drug Name	Indication	Dosing Regimen	Maximum Dose
histrelin acetate (Supprelin® LA)	Central Precocious Puberty	1 implant (50 mg) inserted SC for 12 months	1 implant per 12 months
histrelin acetate (Vantas®)	Advanced prostate cancer palliative therapy	1 implant (50 mg) inserted SC for 12 months	1 implant per 12 months

Dosage Forms

- histrelin acetate (Supprelin® LA): Implant: 50 mg (approximately 65 mcg histrelin acetate per day over 12 months).
- histrelin acetate (Vantas®): Implant: 50 mg (approximately 50 mcg histrelin acetate per day over 12 months).

Clinical Policy

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria. The provision of provider samples does not guarantee coverage under the terms of the pharmacy benefit administered by RxAdvance. All criteria for initial approval must be met in order to obtain coverage.

I. Initial Approval Criteria

A. Prostate Cancer (must meet all):

1. Diagnosis of prostate cancer;
2. Request is for Vantas®;
3. Prescribed by or in consultation with an oncologist or urologist;
4. Age 18 years or older;
5. Request meets one of the following (a or b)*:
 - a. Dose does not exceed 50 mg per 12 months (one implant per year);
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).

This clinical policy has been developed to authorize, modify, or determine coverage for individuals with similar conditions. Specific care and treatment may vary depending on individual need and benefits covered by the plan. This policy is not intended to dictate to providers how to practice medicine, nor does it constitute a contract or guarantee regarding payment or results. This document may contain prescription brand name drugs that are trademarks of pharmaceutical manufacturers that are not affiliated with RxAdvance.

**Prescribed regimen must be FDA-approved or recommended by NCCN.*

Approval Duration

Commercial: 12 months

Medicaid: 12 months

B. Central Precocious Puberty (must meet all):

1. Diagnosis of CPP confirmed by all of the following (a, b, and c):
 - a. Elevated basal luteinizing hormone (LH) level >0.2-0.3 mIU/L (dependent on type of assay used) and or elevated leuprolide-stimulated LH level > 3.3-5 IU/L;
 - b. Difference between bone age and chronological age was > 1 year (bone age chronological age);
 - c. Age at onset of secondary sex characteristics is < 8 years if female, or < 9 years if male;
2. Request is for Supprelin® LA;
3. Prescribed by or in consultation with a pediatric endocrinologist;
4. Member meets one of the following age requirements (a or b):
 - a. Female: 2-11 years;
 - b. Male: 2-12 years;
5. Dose does not exceed 50 mg per 12 months (one implant per year).

Approval Duration

Commercial: 12 months

Medicaid: 12 months

II. Continued Therapy Approval

A. Prostate Cancer (must meet all):

1. Member is currently receiving medication that has been authorized by RxAdvance or the member has met initial approval criteria listed in this policy;
2. Request is for Vantas®;
3. Member is responding positively to therapy;
4. If request is for a dose increase, request meets one of the following (a or b)*:
 - a. New dose does not exceed 50 mg per 12 months (one implant per year);
 - b. New dose is supported by practice guidelines or peer-reviewed literature for relevant off-label use (prescriber must submit supporting evidence).

**Prescribed regimen must be FDA-approved or recommended by NCCN.*

Approval Duration

Commercial: 12 months

Medicaid: 12 months

B. Central Precocious Puberty (must meet all):

1. Member is currently receiving medication that has been authorized by RxAdvance or the member has met initial approval criteria listed in this policy;
2. Request is for Supprelin® LA;
3. Member is responding positively to therapy as evidenced by, including but not limited to, improvement in any of the following parameters: decreased growth velocity, cessation of menses, softening of breast tissue or testes, arrested pubertal progression;
4. Member meets one of the following age requirements (a or b):
 - a. Female: ≤ 11 years;
 - b. Male: ≤ 12 years;
5. If request is for a dose increase, new dose does not exceed 50 mg per 12 months (one implant per year).

Approval Duration

Commercial: 12 months
Medicaid: 12 months

III. Appendices

APPENDIX A: Abbreviation/Acronym Key

CPP: Central Precocious Puberty
FDA: Food and Drug Administration
GnRH: Gonadotropin Releasing Hormone
LH: Luteinizing Hormone
NCCN: National Comprehensive Cancer Network

APPENDIX B: Therapeutic Alternatives

Below are suggested therapeutic alternatives based on clinical guidance. Please check drug formulary for preferred agents and utilization management requirements.

Drug Name	Dosing Regimen	Dose Limit/Maximum Dose
leuprolide acetate injection (generic)	Prostate Cancer - Palliative Therapy SC: 1 mg per day	1 mg per day
Lupron Depot® (leuprolide acetate)	Prostate Cancer - Palliative Therapy IM: 7.5 mg per 4 weeks, 22.5 mg per 12 weeks, 30 mg per 16 weeks, or 45 mg per 24 weeks	See dosing regimen
Eligard® (leuprolide acetate)	Prostate Cancer - Palliative Therapy SC: 7.5 mg per month, 22.5 mg per 3 months, 30 mg per 4 months, or 45 mg per 6 months	See dosing regimen
Zoladex® (goserelin acetate)	Prostate cancer - 10.8 mg SC every 12 weeks	10.8 mg every 12 weeks
Trelstar® (triptorelin pamote)	Prostate cancer – IM: 3.75 mg once every 4 weeks or 11.25 mg once every 12 weeks or 22.5 mg once every 24 weeks	See dosing regimen

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

APPENDIX C: Contraindications/Boxed Warnings

- Contraindication(s):
 - Hypersensitivity to GnRH, GnRH agonist analogs;
 - Pregnancy:
 - Vantas® can cause fetal harm when used during pregnancy.

- Supprelin® LA treatment increases the risk for pregnancy loss.

- Boxed Warning(s):
None.

APPENDIX D: General Information

- **Vantas®:**
 - Transient Increase in Serum Testosterone: Detected during the first week of treatment and may result in worsening of tumor symptoms.
 - Spinal Cord Compression and Urinary Tract Obstruction: May cause paralysis or renal impairment. Monitor patients at risk closely during therapy.
 - Difficulty Locating or Removing Implant: Loss of or inability to locate or remove an inserted implant has been reported. Caution is recommended.
 - Hyperglycemia and Diabetes: Hyperglycemia and an increased risk of developing diabetes have been reported in men receiving GnRH analogs. Monitor blood glucose level and manage according to current clinical practice.
 - Cardiovascular Diseases: Increased risk of myocardial infarction, sudden cardiac death and stroke has been reported in men. Monitor for cardiovascular disease and manage according to current clinical practice.
 - Effect on QT/QTc Interval: Androgen deprivation therapy may prolong the QT interval. Consider risks and benefits.
 - Embryo-Fetal Toxicity: Can cause fetal harm.
- **Supprelin® LA:**
 - Initial Agonistic Action: Initial transient increases of estradiol and/or testosterone may cause a temporary worsening of symptoms.
 - Psychiatric Events: Have been reported in patients taking GnRH agonists. Events include emotional lability, such as crying, irritability, impatience, anger, and aggression. Monitor for development or worsening of psychiatric symptoms.
 - Convulsions have been observed in patients receiving GnRH agonists with or without a history of seizures, epilepsy, cerebrovascular disorders, central nervous system anomalies or tumors, and in patients on concomitant medications that have been associated with convulsions.

References

1. Vantas® Prescribing Information. Malvern, PA: Endo Pharmaceuticals Solutions, Inc.; December 2020. Available at https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/021732s023lbl.pdf. Accessed May 11, 2021.
2. Supprelin® LA Prescribing Information. Malvern, PA: Endo Pharmaceuticals Solutions, Inc.; November 2019. Available at www.supprelinla.com. Accessed May 11, 2021.
3. National Comprehensive Cancer Network Drugs and Biologics Compendium. Histrelin acetate. Available at nccn.org. Accessed May 11, 2021.
4. National Comprehensive Cancer Network. Prostate Cancer Version 2.2021. Available at https://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf. Accessed May 11, 2021.
5. Kaplowitz P, Bloch C. Evaluation and referral of children with signs of early puberty. *Pediatrics*. 2016; 137(1): e20153732.
6. Vantas & Supprelin LA Prescribing Information. Endo Pharmaceuticals Inc. Available at <https://dailymed.nlm.nih.gov/dailymed/druginfo.cfm?setid=d8fb000e-3cc9-4803-b71d-2cc597661977> December 13, 2019. Accessed June 25, 2020.
7. Goserelin acetate; In: UpToDate, Post, TW (Ed), UpToDate, Waltham, MA, 2021. Accessed with subscription at: <http://uptodate.com>. Accessed May 11, 2021.
8. Triptorelin pamoate; In: UpToDate, Post, TW (Ed), UpToDate, Waltham, MA, 2021. Accessed with subscription at:

<http://uptodate.com>. Accessed May 11, 2021.

9. Leuprolide acetate; In: UpToDate, Post, TW (Ed), UpToDate, Waltham, MA, 2021. Accessed with subscription at: <http://uptodate.com>. Accessed May 11, 2021.
10. Eugster EA. Treatment of Central Precocious Puberty. J Endocr Soc. 2019;3(5):965-972. Published 2019 Mar 28. doi:10.1210/js.2019-00036.
11. Klein DA, Emerick JE, Sylvester JE, Vogt KS. Disorders of Puberty: An Approach to Diagnosis and Management. American Family Physician. 2017 Nov;96(9):590-599.

Review/Revision History	Review/Revised Date	P&T Approval Date
Policy established.	01/2020	02/07/2020
Policy was reviewed: <ol style="list-style-type: none"> 1) Policy description table was updated. 2) Continued therapy criteria II.A.1 and II.B.1 was rephrased to “Member is currently receiving medication that has been authorized by RxAdvance...”. 3) Initial therapy and continued therapy criteria were updated to include commercial and Medicaid separately. 4) Appendix B, therapeutic alternatives table was added. Alternative drugs were listed per NCCN prostate cancer guidelines. 5) Appendix C, contraindications was updated. 6) References updated. 	06/25/2020	09/14/2020
Policy was reviewed: <ol style="list-style-type: none"> 1. Policy title table updated. 2. Clinical policy section standard verbiage was updated to include “The provision of prescriber samples...”. 3. Continued therapy approval criteria II.A.1 was rephrased to “Member is currently receiving medication...”. 4. Appendix B for therapeutic alternatives standard verbiage was updated to “Below are suggested therapeutic alternatives based on clinical guidance...”. 5. Appendix D for general information was added. 6. References were updated. 	05/11/2021	06/10/2021